


AMENDMENTS TO THE CLAIMS

1. (Currently amended) A cinnamoyl compound represented by the formula (I):



wherein:


~~I. (I)~~ A represents a ~~benzene-phenyl~~ ring or a ~~pyridine-pyridyl~~ ring; and in $(Y_\alpha)_q$, Y_α is a substituent on a carbon atom and represents a group included in the following X_0 group or Y_0 group, q represents 0, 1, 2, 3 or 4, and Y_α s are the same or different when q is 2 or more and the adjacent two same or different Y_α s together may form a group included in the Z_0 group to be fused to the A ring when q is 2 or more; and in $(X_\alpha)_p$, X_α ~~represents-is~~ a substituent on a carbon atom and represents a_{3-1} -CH₂-CO-NH- (wherein a_{3-1} represents a C1-C10 alkoxy group) ~~which does not belong to the following X_0 group, Y_0 group and Z_0 group~~, p represents 1, 2, 3, 4 or 5, and X_α s may be the same or different when p is 2 or more; and the sum of p and q is 5 or less;

(1) the X_0 group: a M_a -group, wherein M_a represents a R_b - group (wherein R_b represents a C1-C10 alkyl group optionally substituted with a halogen atom), a halogen atom, a nitro group, a cyano group, a hydroxy group, a R_c - B_a - R_d - group (wherein R_c represents a C1-C10 alkyl group optionally substituted with a halogen atom, B_a represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group, and R_d represents a single bond or a C1-C10 alkylene group), a HOR_d - group (wherein R_d is as defined above), a R_c -CO- R_d - group (wherein R_c represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom, and R_d is as defined above), a R_c -CO-O- R_d - group (wherein R_c and R_d are as defined above), a R_c O-CO- R_d - group (wherein R_c and R_d are as defined above), a HO-CO-CH=CH- group, a R_cR_c' -N- R_d - group (wherein R_c and R_c' are the same or different, R_c is as defined above, R_c' has the same meaning as R_c has, and R_d is as defined above), a R_c -CO-N R_c' - R_d - group (wherein R_c , R_c' and R_d are as defined above), a R_b O-CO-N(R_c)- R_d - group (wherein R_b , R_c and R_d are as defined above), a R_cR_c' -N-CO- R_d - group (wherein R_c , R_c' and R_d are as defined above), a R_cR_c' -N-CO-N R_c'' - R_d - group (wherein R_c , R_c' and R_c'' are the same or different, R_c and R_c' are as defined above, R_c''

has the same meaning as R_e has, and R_d is as defined above), a $R_e R_e' N-C(=NR_e'')-NR_e'''-R_d$ -group (wherein R_e , R_e' , R_e'' and R_e''' are the same or different, R_e , R_e' and R_e'' are as defined above, R_e''' has the same meaning as R_e has, and R_d is as defined above), a $R_b-SO_2-NR_e-R_d$ -group (wherein R_b , R_e and R_d are as defined above), a $R_e R_e' N-SO_2-R_d$ -group (wherein R_e , R_e' and R_d are as defined above), a C2-C10 alkenyl group or a C2-C10 alkynyl group;

(2) the Y_0 group: a $M_{b0}-R_d$ -group, wherein M_{b0} represents a M_{c0} -group


~~{wherein~~ wherein M_{c0} represents a $M_{d0}-R_d'$ -group ~~{wherein~~ wherein M_{d0} represents a 6 to 10-membered aryl group optionally substituted with a M_a -group (wherein M_a is as defined above), a 5 to 10-membered heteroaryl group optionally substituted with a M_a -group (wherein M_a is as defined above), a 3 to 10-membered cyclic hydrocarbon or heterocyclic group optionally substituted with a M_a -group (wherein M_a is as defined above) and optionally containing an unsaturated bond, a (b_0) -group




(in the (b_0) -group, G_0 forms an optionally substituted, saturated or unsaturated, nonaromatic 5 to 14-membered cyclic hydrocarbon or heterocyclic ring), a (c_0) -group



(in the (c_0) -group, J_0 forms a 5 to 7-membered aromatic ring optionally containing a nitrogen atom), a (d_0) -group



~~{wherein~~ wherein d_0 forms a 5 to 12-membered hydrocarbon ring which is substituted with a carbonyl group or a thiocarbonyl group and further which may be optionally substituted with an oxy group, a thio group, a $-NR_1$ -group {wherein R_1 represents a hydrogen atom, a C1-C10 alkyl group, a C2-C10 alkyl group substituted with a halogen atom or a R_2-B_1 -group (wherein R_2 represents a C1-C10 alkyl group, a C3-C10 alkenyl group or a C3-C10 alkynyl group, and B_1 represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group), a C3-C10 alkenyl group, or a C3-C10 alkynyl group}, a sulfinyl group or a sulfonyl group} or a (e_0) -group



{wherein e₀ forms a 5 to 12-membered hydrocarbon ring optionally substituted with a carbonyl group, a thiocarbonyl group, an oxy group, a thio group, a -NR₁- group (wherein R₁ is as defined above), a sulfinyl group or a sulfonyl group}; and R_d' is the same as or different from R_d and has the same meaning as R_d ~~has~~ has),

a M_{c0}-B_a- group (wherein M_{c0} and B_a are as defined above), a M_{c0}-CO- group (wherein M_{c0} is as defined above), a M_{c0}-CO-O- group (wherein M_{c0} is as defined above), a M_{c0}O-CO- group (wherein M_{c0} is as defined above), a M_{c0}R_eN- group (wherein M_{c0} and R_e are as defined above), a M_{c0}-CO-NR_e- group (wherein M_{c0} and R_e are as defined above), a M_{c0}O-CO-NR_e- group (wherein M_{c0} and R_e are as defined above), a M_{c0}R_eN-CO- group (wherein M_{c0} and R_e are as defined above), a M_{c0}R_eN-CO-NR_e'- group (wherein M_{c0}, R_e and R_e' are as defined above), a M_{c0}R_eN-C(=NR_e')-NR_e''- group (wherein M_{c0}, R_e, R_e' and R_e'' are as defined above), a M_{c0}-SO₂-NR_e- group (wherein M_{c0} and R_e are as defined above) or a M_{c0}R_eN-SO₂- group (wherein M_{c0} and R_e are as defined above), and

R_d is as defined above;

(3) the Z₀ group: a 5 to 12-membered cyclic hydrocarbon or heterocyclic ring optionally substituted with a halogen atom, a C1-C10 alkoxy group, a C3-C10 alkenyloxy group, a C3-C10 alkynyloxy group, a carbonyl group, a thiocarbonyl group, an oxy group, a thio group, a sulfinyl group or a sulfonyl group, which is an aromatic or nonaromatic and monocyclic or fused ring and which is fused to the A ring;


~~H-(II)~~ Q_a represents an optionally substituted hydroxy group, or an optionally substituted amino group;

~~HH-(III)~~ K_a and L_a are the same or different, and represent a hydrogen atom, or a substituent on a carbon atom, or K_a and L_a may form a C1-C10 alkylene group optionally having a substituent or a C1-C10 alkenylene group optionally having a substituent; and

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the

same, selected substituents may be the same or different as long as they are selected within the range;range.

2. (Currently amended) A cinnamoyl compound represented by the formula (II):



wherein:

I-(I) A represents a benzene-phenyl ring or a pyridine-pyridyl ring;

II-(II) in $(X_{A0})_p$, X_{A0} is a substituent on a carbon atom and represents $a_{3-1}-CH_2-CO-NH-$ (wherein a_{3-1} represents a C1-C10 alkoxy group) and ~~represents a group included in any group of the following A_0 to N_0 groups,~~ p represents 1, 2, 3, 4 or 5, and when p is 2 or more, X_{A0} s are the same or different;

(1) the A_0 group:

~~—— a D_1-R_4 group [wherein D_1 represents a $(R_1-(O)_k)-A_1N-(O)_k$ group [wherein R_1 represents a hydrogen atom, or a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a halogen atom or a R_2-B_1 group (wherein R_2 represents a C1-C10 alkyl group, a C3-C10 alkenyl group or a C3-C10 alkynyl group, and B_1 represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group), or a C3-C10 alkenyl group, or a C3-C10 alkynyl group, k represents 0 or 1, A_1 represents a $R_2-(CHR_0)_m-(B_2-B_3)_m$ group (wherein R_2 represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom or a R_2-B_1 group (wherein R_2 and B_1 are as defined above), or a C2-C10 alkenyl group, or a C2-C10 alkynyl group, R_0 represents a hydrogen atom, a C1-C10 alkyl group or a C2-C10 haloalkyl group, m represents 0 or 1, B_2 represents a single bond, an oxy group, a thio group or a $N((O)_nR_4')$ group (wherein R_4' is the same as or different from R_4 , and has the same meaning as R_4 has, and n represents 0 or 1), B_3 represents a carbonyl group, a thiocarbonyl group or a sulfonyl group, m' represents 0 or 1, and when B_3 is a sulfonyl group, it does not occur that m is 0 and R_4 is a hydrogen atom at the same~~


time}, and k' represents 0 or 1}, and R₄ represents a C1-C10 alkylene group, provided that a R₀'R₀'N-R₄ group (wherein R₀' and R₀' are the same as or different from R₀ and have the same meaning as R₀ has, and R₄ is as defined above) is excluded],

— a D₂-R₄ group [wherein D₂ represents a cyano group, a R₁R₁'NC(=N(O)_n-A₁) group (wherein R₁, R₁', n and A₁ are as defined above), an A₁N=C(OR₂) group (wherein A₁ and R₂ are as defined above) or a NH₂-CS group, and R₄ is as defined above],

— a D₃-R₄ group [wherein D₃ represents a nitro group or a R₁OSO₂ group (wherein R₁ is as defined above), and R₄ is as defined above], or

— a R₁OSO₂ group [wherein R₁ is as defined above];


— (2) the B₀ group: an (a₀) group




in the (a₀) group, E₀ forms an optionally substituted, saturated or unsaturated, aromatic or nonaromatic 5 to 14-membered cyclic hydrocarbon or heterocyclic ring, and R₁ is as defined above;

— (3) the C₀ group: a C2-C10 alkenyl group substituted with a halogen atom, a R₂-B₁ group (wherein R₂ and B₁ are as defined above), a D₄-R₄ group [wherein D₄ represents a hydroxy group or an A₁-O group (wherein A₁ is as defined above), and R₄ is as defined above], a D₅ group [wherein D₅ represents a O=C(R₂) group (wherein R₂ is as defined above), an A₁-(O)_n-N=C(R₂) group (wherein A₁, n and R₂ are as defined above), a R₁-B₀-CO-R₄-(O)_n-N=C(R₂) group (wherein R₁, R₄, n and R₂ are as defined above, and B₀ represents an oxy group, a thio group or a N((O)_mR₁') group (wherein R₁' and m are as defined above)], a D₂-R₄-(O)_n-N=C(R₂) group (wherein D₂, R₄, n and R₂ are as defined above) or a R₁A₁N-N=C(R₂) group (wherein R₁, A₁ and R₂ are as defined above)], a R₁A₁N-O-R₄ group (wherein R₁, A₁ and R₄ are as defined above), a R₁(A₁-(O)_n)N group (wherein R₁, A₁ and n are as defined above), a D₂ group (wherein D₂ is as defined above) or a D₃ group (wherein D₃ is as defined above);

— (4) the D₀ group: a C2-C10 alkynyl group substituted with a (b₀)-R₄ group (in (b₀))



G_0 forms an optionally substituted, saturated or unsaturated, nonaromatic 5 to 14 membered cyclic hydrocarbon or heterocyclic ring), a (e_0) - R_4 -group (in (e_0)



J_0 forms an aromatic 5 to 7 membered ring optionally containing a nitrogen atom and R_4 is as defined above), a halogen atom, a R_2 - B_4 - R_4 -group (wherein R_2 , B_4 and R_4 are as defined above), a D_4 - R_4 -group (wherein D_4 and R_4 are as defined above), a D_5 -group (wherein D_5 is as defined above), a D_1 - R_4 -group (wherein D_1 and R_4 are as defined above), a D_2 -group (wherein D_2 is as defined above) or a D_3 - R_4 -group (wherein D_3 and R_4 are as defined above);

—— (5) the E_0 group: an A_2 -CO- R_5 -group, provided that R_5 is not a vinylene group when A_2 is a hydroxy group, wherein A_2 represents

—— (i) an A_3 - B_4 -group

—— wherein A_3 represents a hydrogen atom, or a C1-C10 alkyl group, or a C2-C10 haloalkyl group, or a C2-C10 alkenyl group optionally substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or a R_{a0} -(R_4)_m-group (wherein R_{a0} represents an optionally substituted 5 to 7 membered aryl group or heteroaryl group, and R_4 and m are as defined above), or a C1-C10 alkyl group substituted with a (b_0) - R_4 -group (wherein (b_0) and R_4 are as defined above), a (e_0) - R_4 -group (wherein (e_0) and R_4 are as defined above), a R_2 - B_4 - R_4 -group (wherein R_2 , B_4 and R_4 are as defined above), a D_4 - R_4 -group (wherein D_4 and R_4 are as defined above), a D_5 -group (wherein D_5 is as defined above), a D_1 - R_4 -group (wherein D_1 and R_4 are as defined above), a D_2 -group (wherein D_2 is as defined above), a D_3 - R_4 -group (wherein D_3 and R_4 are as defined above) or an A_4 -SO₂- R_4 -group (wherein A_4 represents a (b_0) -group (wherein (b_0) is as defined above), a (e_0) -group (wherein (e_0) is as defined above) or a R_1R_1' -N-group (wherein R_1 and R_1' are as defined above), and R_4 is as defined above), and

—— B_4 represents an oxy group, a thio group or a N((O)_m R_1)-group (wherein R_1 and m are as defined above), provided that A_2 is not a hydrogen atom when B_4 is a thio group;

—— (ii) a R_1 - B_4 -CO- R_4 - B_4' -group, wherein R_1 , B_4 and R_4 are as defined above, B_4' is the same as or different from B_4 and has the same meaning as B_4 has, provided that R_2 is not a hydrogen atom when B_4 is a thio group, or

—— a D_2 - R_4 - B_4 -group, wherein D_2 , R_4 and B_4 are as defined above;

~~—— (iii) a $R_2-SO_2-NR_4$ group, wherein R_2 is as defined above, provided that a hydrogen atom is excluded, and R_4 is as defined above;~~

~~—— (iv) a (b_0) group, wherein (b_0) is as defined above;~~

~~(v) a (e_0) group, wherein (e_0) is as defined above; or~~

~~—— (vi) a $R_4A_4N-NR_4'$ group, wherein R_4 , A_4 and R_4' are as defined above; and~~

~~R_5 represents a C2-C10 alkenylene group optionally substituted with a halogen atom or a C2-C10 alkynylene group;~~

~~—— (6) the F_0 group: an $A_5-B_5-R_6$ group~~

~~—— wherein A_5 represents a C2-C10 alkyl group substituted with a D_4 group (wherein D_4 is as defined above), a D_1 group (wherein D_1 is as defined above), a D_3 group (wherein D_3 is as defined above) or an A_4-SO_2 group (wherein A_4 is as defined above), or a C1-C10 alkyl group substituted with a R_2-B_1 group (wherein R_2 and B_1 are as defined above), a D_2 group (wherein D_2 is as defined above), a D_5 group (wherein D_5 is as defined above) or an A_2-CO group (wherein A_2 is as defined above);~~

~~—— B_5 represents a B_1 group (wherein B_1 is as defined above) or a NA_4 group (wherein A_4 is as defined above), and~~

~~—— R_6 represents a single bond or a C1-C10 alkylene group;~~

~~(7) the G_0 group: an $A_6-B_5-R_6$ group~~

~~—— wherein A_6 represents an (a_0) R_4 group (wherein (a_0) and R_4 are as defined above), or a C2-C10 alkenyl group, or a C2-C10 alkynyl group, or a C2-C10 alkenyl group substituted with a halogen atom, a R_2-B_1 group (wherein R_2 and B_1 are as defined above), a D_5 group (wherein D_5 is as defined above), a D_2 group (wherein D_2 is as defined above) or an A_2-CO group (wherein A_2 is as defined above), or a C2-C10 alkynyl group substituted with a halogen atom, a R_2-B_1 group (wherein R_2 and B_1 are as defined above), a D_5 group (wherein D_5 is as defined above), a D_2 group (wherein D_2 is as defined above) or an A_2-CO group (wherein A_2 is as defined above), or a C3-C10 alkenyl group substituted with a (b_0) group (wherein (b_0) is as defined above), a (e_0) group (wherein (e_0) is as defined above), a D_4 group (wherein D_4 is as defined above), a D_1 group (wherein D_1 is as defined above) or a D_3 group (wherein D_3 is as defined above), or a C3-C10 alkynyl group substituted with a D_4 group (wherein D_4 is as defined~~

above), a D_4 -group (wherein D_4 is as defined above) or a D_3 -group (wherein D_3 is as defined above), and

—— B_5 and R_6 are as defined above;

—— (8) the H_0 group:

—— a $D_2-N((O)_nA_4)-R_6$ -group (wherein D_2 , n , A_4 and R_6 are as defined above);

—— a D_2 -group (wherein D_2 is as defined above, provided that a cyano group is excluded);

—— a $R_4(R_4'(O)_n)N-CR_4''-N-R_6$ -group (wherein R_4 , R_4' , n and R_6 are as defined above, R_4'' is the same as or different from R_4 and has the same meaning as that of R_4);

—— a $R_4-(O)_n-N-CR_4'-NR_2-R_6$ -group (wherein R_4 , n , R_4' , R_2 and R_6 are as defined above);

—— a $R_2-B_3-NR_4-CO-NR_4'-R_6$ -group (wherein R_2 , B_3 , R_4 , R_4' and R_6 are as defined above);

—— a $D_2-CO-NR_4-R_6$ -group (wherein D_2 , R_4 and R_6 are as defined above) or

—— an $A_3-COCO-NR_4-R_6$ -group (wherein A_3 , R_4 and R_6 are as defined above);

—— (9) the I_0 group:

—— an $A_7-B_6-N((O)_nR_4)-R_6$ -group [wherein A_7 represents a C2-C10 alkenyl group optionally substituted with a halogen atom, or a C2-C10 alkynyl group, or a C3-C10 haloalkynyl group, or a $R_2-B_4-R_4$ -group (wherein R_2 , B_4 and R_4 are as defined above), or a D_4-R_4 -group (wherein D_4 and R_4 are as defined above), or a D_5-R_4 -group (wherein D_5 and R_4 are as defined above), or a D_1-R_4 -group (wherein D_1 and R_4 are as defined above), or a $(b_0)-R_4$ -group (wherein (b_0) and R_4 are as defined above), or a $(c_0)-R_4$ -group (wherein (c_0) and R_4 are as defined above), or a D_2-R_4 -group (wherein D_2 and R_4 are as defined above), or a D_3-R_4 -group (wherein D_3 and R_4 are as defined above), or an $A_4-SO_2-R_4$ -group (wherein A_4 and R_4 are as defined above), or an A_2-CO-R_4 -group (wherein A_2 and R_4 are as defined above), B_6 represents a carbonyl group or a thiocarbonyl group, and n , R_4 and R_6 are as defined above];

—— an $A_8-CS-N((O)_nR_4)-R_6$ -group [wherein A_8 represents a hydrogen atom or a C1-C10 alkyl group optionally substituted with a halogen atom, and n , R_4 and R_6 are as defined above];

—— an $A_7'-B_2'-B_3-N((O)_nR_4)-R_6$ -group [wherein A_7' represents a C3-C10 alkenyl group optionally substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or a $R_2-B_4-R_4'$ -group (wherein R_2 and B_4 are as defined above, and R_4' represents a C2-C10 alkylene group), or a D_4-R_4' -group (wherein D_4 and R_4' are as defined above), or a D_1-R_4' -group (wherein D_1 and R_4' are as defined above), or a $(b_0)-R_4'$ -group (wherein (b_0) and R_4' are as defined above), or a $(c_0)-R_4'$ -group (wherein (c_0) and R_4' are as

defined above), or a D_2-R_4 group (wherein D_2 and R_4 are as defined above), or a D_3-R_4' group (wherein D_3 and R_4' are as defined above), or an A_2-CO-R_4 group (wherein A_2 and R_4 are as defined above), B_2' represents an oxy group, a thio group or a $N((O)_nR_4')$ group (wherein n' is the same as or different from n and has the same meaning as that of n , and R_4' is as defined above), and B_3 , n , R_4 and R_6 are as defined above];

—— an $A_8'-B_2'-CS-N((O)_nR_4)-R_6$ group [wherein A_8' represents a C1-C10 alkyl group or a C2-C10 haloalkyl group, B_2' is as defined above, and n , R_4 and R_6 are as defined above];

—— an $A_8'-S-B_2'-N((O)_nR_4)-R_6$ group [wherein A_8' , n , R_4 and R_6 are as defined above, and B_2' represents a carbonyl group or a sulfonyl group] or

—— an $A_7''-SO_2-N((O)_nR_4)-R_6$ group [wherein A_7'' represents a C2-C10 alkenyl group, or a C3-C10 alkenyl group substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or a $R_2-B_4-R_4'$ group (wherein R_2 , B_4 and R_4' are as defined above), or a D_4-R_4' group (wherein D_4 and R_4' are as defined above), or a D_5-R_4 group (wherein D_5 and R_4 are as defined above), or a D_1-R_4' group (wherein D_1 and R_4' are as defined above), or a $(b_0)-R_4'$ group (wherein (b_0) and R_4' are as defined above), or a $(e_0)-R_4'$ group (wherein (e_0) and R_4' are as defined above), or a D_2-R_4 group (wherein D_2 and R_4 are as defined above), or a NO_2-R_4 group (wherein R_4 is as defined above), or an A_2-CO-R_4 group (wherein A_2 and R_4 are as defined above), and n , R_4 and R_6 are as defined above];

—— (10) the J_0 group:

—— an A_7-CO group (wherein A_7 is as defined above),

—— an A_9-CS group (wherein A_9 represents A_7 or A_8),

—— an $A_9'(O)_mN-C(A_9)$ group (wherein A_9' represents A_7' or A_8' , and m and A_9 are as defined above),

—— a D_2-CO group (wherein D_2 is as defined above),

—— an A_2-COCO group (wherein A_2 is as defined above),

—— an $A_9-CO-B_4'-R_6$ group (wherein A_9 and R_6 are as defined above, and B_4' represents an oxy group or a thio group, provided that A_9 is not A_8 when B_4' is an oxy group),

—— an $A_9-CS-B_4'-R_6$ group (wherein A_9 , B_4' and R_6 are as defined above),

—— an $A_7''-SO_2-B_4'-R_6$ group (wherein A_7'' , B_4' and R_6 are as defined above),

—— an $A_8-SO_2-B_4'-R_6$ group (wherein A_8 , B_4' and R_6 are as defined above, provided that A_8 is not a hydrogen atom),

~~— an A_9' - B_2' - B_2 - B_4' - R_6 group (wherein A_9' , B_2' , B_2 , B_4' and R_6 are as defined above), or~~
~~— a C2-C10 alkenyl group substituted with a (b_0) group (wherein (b_0) is as defined above)~~
~~or a (c_0) group (wherein (c_0) is as defined above);~~

~~— (11) the K_0 group: an A_{10} - $N((O)_nR_1)$ -CO- R_6 group~~
~~— wherein A_{10} represents a hydrogen atom (provided that n is not 0), an A_2'' -SO₂ group (wherein A_2'' is as defined above), an A_8 -SO₂ group (wherein A_8 is as defined above, provided that A_8 is not a hydrogen atom), an A_9' -O group (wherein A_9' is as defined above, provided that n is not 1), an A_9' group (wherein A_9' is as defined above, provided that A_8' is excluded when n is 0), a R_2 OCH₂ group (wherein R_2 is as defined above), an A_2 -CO- R_4 group (wherein A_2 and R_4 are as defined above) or an A_2 -CO-CH(CH₂CO- A_2) group (wherein A_2 is as defined above), and n, R_1 and R_6 are as defined above;~~

~~— (12) the L_0 group:~~
~~— an A_{10}' - $N((O)_nR_1)$ -SO₂- R_6 group [wherein A_{10}' represents a hydrogen atom (provided that n is not 0), an A_9' -O group (wherein A_9' is as defined above, provided that n is not 1), an A_9' group (wherein A_9' is as defined above, provided that A_8' is excluded when n is 0), a R_2 -CO group (wherein R_2 is as defined above), an A_2 -CO- R_4 group (wherein A_2 and R_4 are as defined above) or an A_2 -CO-CH(CH₂CO- A_2) group (wherein A_2 is as defined above), and n, R_1 and R_6 are as defined above];~~

~~— an A_9'' - R_1 N-SO₂- $N((O)_nR_1')$ - R_6 group [wherein A_9'' represents a hydrogen atom or an A_9' group (wherein A_9' is as defined above), and R_1 , n, R_1' and R_6 are as defined above] or~~
~~— a (b_0) -SO₂- $N((O)_nR_1')$ - R_6 group [wherein (b_0) , n, R_1' and R_6 are as defined above];~~

~~— (13) the M_0 group:~~
~~— a R_1 (R_2 S)C=N- R_6 group (wherein R_1 , R_2 and R_6 are as defined above),~~
~~— a R_2 B(R_2' B')C=N- R_6 group (wherein R_2 and R_6 are as defined above, R_2' is the same as or different from R_2 and has the same meaning as that of R_2 , and B and B' are the same or different and represent an oxy group or a thio group),~~

~~— a R_1 R_1' N(R_2 S)C=N- R_6 group (wherein R_1 , R_1' , R_2 and R_6 are as defined above),~~
~~— a R_1 N=C(S R_2)NR₂'- R_6 group (wherein R_1 , R_2 , R_2' and R_6 are as defined above) or~~
~~— a R_1 (R_1' O)N- R_6 group (wherein R_1 , R_1' and R_6 are as defined above);~~

~~— (14) the N_0 group: a A_{11} -P(=O)(OR₁')- R_4 group~~

~~wherein A_{11} represents a R_1 -group (wherein R_1 is as defined above), a R_1O-R_6 -group (wherein R_1 and R_6 are as defined above) or a $R_1OCO-CHR_6$ -group (wherein R_1 and R_6 are as defined above), and R_1' and R_4 are as defined above;~~


III.(III) in $(Y_{\Lambda 0})_q$, $Y_{\Lambda 0}$ is a substituent on a carbon atom and represents a group included in the following X_0 group and Y_0 group, q represents 0, 1, 2, 3 or 4, the sum of p (wherein p is as defined above) and q is 5 or less, $Y_{\Lambda 0}$ s are the same as or different when q is 2 or more, and the adjacent two same or different $Y_{\Lambda 0}$ s may form a group included in the Z_0 group to be fused to the A ring when q is 2 or more;

(1) the X_0 group: a M_a - group, wherein M_a represents a R_b - group (wherein R_b represents a C1-C10 alkyl group optionally substituted with a halogen atom), a halogen atom, a nitro group, a cyano group, a hydroxy group, a $R_c-B_a-R_d$ - group (wherein R_c represents a C1-C10 alkyl group optionally substituted with a halogen atom, B_a represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group, and R_d represents a single bond or a C1-C10 alkylene group), a HOR_d - group (wherein R_d is as defined above), a R_c-CO-R_d - group (wherein R_c represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom, and R_d is as defined above), a $R_c-CO-O-R_d$ - group (wherein R_c and R_d are as defined above), a $R_cO-CO-R_d$ - group (wherein R_c and R_d are as defined above), a $HO-CO-CH=CH$ - group, a $R_cR_c'N-R_d$ - group (wherein R_c and R_c' are the same or different, R_c is as defined above, R_c' has the same meaning as R_c has, and R_d is as defined above), a $R_c-CO-NR_c'-R_d$ - group (wherein R_c , R_c' and R_d are as defined above), a $R_bO-CO-N(R_c)-R_d$ - group (wherein R_b , R_c and R_d are as defined above), a $R_cR_c'N-CO-R_d$ - group (wherein R_c , R_c' and R_d are as defined above), a $R_cR_c'N-CO-NR_c''-R_d$ - group (wherein R_c , R_c' and R_c'' are the same or different, R_c and R_c' are as defined above, R_c'' has the same meaning as R_c has, and R_d is as defined above), a $R_cR_c'N-C(=NR_c'')-NR_c'''-R_d$ - group (wherein R_c , R_c' , R_c'' and R_c''' are the same or different, R_c , R_c' and R_c'' are as defined above, R_c''' has the same meaning as R_c has, and R_d is as defined above), a $R_b-SO_2-NR_c-R_d$ - group (wherein R_b , R_c and R_d are as defined above), a $R_cR_c'N-SO_2-R_d$ - group (wherein R_c , R_c' and R_d are as defined above), a C2-C10 alkenyl group or a C2-C10 alkynyl group;


(2) the Y_0 group: a $M_{b0}-R_d$ - group, wherein M_{b0} represents a M_{c0} - group

~~wherein~~ (wherein M_{c0} represents a $M_{d0}-R_d$ - group ~~wherein~~ (wherein M_{d0} represents a 6 to 10-membered aryl group optionally substituted with a M_a - group (wherein M_a is as defined


above), a 5 to 10-membered heteroaryl group optionally substituted with a M_a - group (wherein M_a is as defined above), a 3 to 10-membered cyclic hydrocarbon or heterocyclic group which is optionally substituted with a M_a - group (wherein M_a is as defined above) and which optionally contains an unsaturated bond, or a (b_o) - group




(wherein (b_o) forms as defined above), a (c_o) - group



(wherein (c_o) forms as defined above), a (d_o) - group



{wherein d_o forms a 5 to 12-membered hydrocarbon ring which is substituted with a carbonyl group or a thiocarbonyl group and further which may be optionally substituted with an oxy group, a thio group, a $-NR_1$ - group (wherein R_1 is a hydrogen atom, or a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a halogen atom or a R_2 - B_1 -group (wherein R_2 represents a C1-C10 alkyl group, a C3-C10 alkenyl group or a C3-C10 alkynyl group, and B_1 represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group), or a C3-C10 alkenyl group, or a C3-C10 alkynyl group as defined above), a sulfinyl group or a sulfonyl group} or a (e_o) - group



{wherein e_o forms a 5 to 12-membered hydrocarbon ring optionally substituted with a carbonyl group, a thiocarbonyl group, an oxy group, a thio group, a $-NR_1$ - group (wherein R_1 is as defined above), a sulfinyl group or a sulfonyl group}, and R_d' is the same as or different from R_d and has the same meaning as R_d ~~has~~ has),

a M_{c0} - B_a - group (wherein M_{c0} and B_a are as defined above), a M_{c0} -CO- group (wherein M_{c0} is as defined above), a M_{c0} -CO-O- group (wherein M_{c0} is as defined above), a M_{c0} O-CO- group (wherein M_{c0} is as defined above), a $M_{c0}R_eN$ - group (wherein M_{c0} and R_e are as defined above), a M_{c0} -CO-NR_e- group (wherein M_{c0} and ~~Re~~ R_e are as defined above), a M_{c0} O-CO-NR_e- group

(wherein M_{c0} and R_c are as defined above), a $M_{c0}R_cN-CO-$ group (wherein M_{c0} and R_c are as defined above), a $M_{c0}R_cN-CO-NR_c'-$ group (wherein M_{c0} , R_c and R_c' are as defined above), a $M_{c0}R_cN-C(=NR_c')-NR_c''-$ group (wherein M_{c0} , R_c , R_c' and R_c'' are as defined above), a $M_{c0}-SO_2-NR_c-$ group (wherein M_{c0} and R_c are as defined above) or a $M_{c0}R_cN-SO_2-$ group (wherein M_{c0} and R_c are as defined above), and R_d is as defined above;

(3) the Z_0 group: a 5 to 12-membered cyclic hydrocarbon or heterocyclic ring optionally substituted with a halogen atom, a C1-C10 alkoxy group, a C3-C10 alkenyloxy group, a C3-C10 alkynyloxy group, a carbonyl group, a thiocarbonyl group, an oxy group, a thio group, a sulfinyl group or a sulfonyl group, which is an aromatic or nonaromatic and monocyclic or fused ring and which is fused to the A ring;


~~IV.~~(IV) Q_{A0} represents a hydroxyl group, a $(b_0)-$ group (wherein (b_0) is as defined above), an $A_9-B_6-B_c-$ group ~~{wherein~~(wherein A_9 and B_6 are as defined above, and B_c represent an oxy group or a $-N((O)_mR_1)-$ group (wherein m and R_1 are as defined above), provided that B_c is not a sulfonyl group when A_9 is a hydrogen atom~~atom~~), an $A_7''-SO_2-B_c-$ group (wherein A_7'' and B_c are as defined above), an $A_8-SO_2-B_c-$ group (wherein A_8 and B_c are as defined above, provided that A_8 is not a hydrogen atom), a $R_1R_1'N-SO_2-B_c-$ group (wherein R_1 , R_1' and B_c are as defined above), a $(b_0)-SO_2-B_c-$ group (wherein (b_0) and B_c are as defined above), an $A_9'-B_c-$ group (wherein A_9' and B_c are as defined above), a $D_5-R_4-B_c-$ group (wherein D_5 , R_4 and B_c are as defined above), a $M_{c0}-B_3-B_c-$ group (wherein M_{c0} , B_3 and B_c are as defined above) or a $M_{c0}-B_c-$ group (wherein M_{c0} and B_c are as defined above);

~~V.~~(V) K_{A0} represents a hydrogen atom, a halogen atom, or a C10 alkyl group, L_{A0} represents a hydrogen atom, or a M_{b0} -group (M_{b0} is as defined above), or K_{A0} and L_{A0} may form a C1-C10 alkylene group, or a C1-C10 alkenylene group optionally substituted with single or the same or different plural M_a groups; and

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the ~~range;~~range.

3. (Currently amended) A cinnamoyl compound represented by the formula (III):

(III)



wherein:

I.(I) A represents a benzene-phenyl ring or a pyridine-pyridyl ring;

II. in $(X_A)_p$, X_A is a substituent on a carbon atom and represents a_{3-1} -CH₂-CO-NH- (wherein a_{3-1} represents a C1-C10 alkoxy group) and a group included in any group or the following A to N groups, p represents 1, 2, 3, 4 or 5, and; X_A s are the same or different when p is 2 or more;

(1) the A group:


~~— a D₁-R₄-group, wherein D₁ represents a (R₄-(O)_k-(A₁N-(O)_k)' group [wherein R₄ represents a hydrogen atom, or a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a halogen atom or a R₂-B₁-group (wherein R₂ represents a C1-C10 alkyl group, a C3-C10 alkenyl group or a C3-C10 alkynyl group, and B₁ represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group), or a C3-C10 alkenyl group, or a C3-C10 alkynyl group, k represents 0 or 1, A₁ represents a R₂-(CHR₀)_m-(B₂-B₂)_{m'}-group (wherein R₂ represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom or a R₂-B₁-group (wherein R₂ and B₁ are as defined above), or a C2-C10 alkenyl group, or a C2-C10 alkynyl group, R₀ represents a hydrogen atom, a C1-C10 alkyl group or a C2-C10 haloalkyl group, m represents 0 or 1, B₂ represents a single bond, an oxy group, a thio group or a N((O)_nR₄') group (wherein R₄' is the same as or different from R₄ and has the same meaning as R₄ has, and n represents 0 or 1), B₂ represents a carbonyl group, a thiocarbonyl group or a sulfonyl group, m' represents 0 or 1, and when B₂ is a sulfonyl group, it does not occur that m is 0 and R₂ is a hydrogen atom at the same time), and k' represents 0 or 1], and R₄ represents a C1-C10 alkylene group, provided that a R₀'R₀'N-R₄-group (wherein R₀' and R₀' are the same as or different from R₀ and has the same meaning as R₀ has, and R₄ is as defined above) is excluded,~~

~~—— a D_2 - R_4 group, wherein D_2 represents a cyano group, a $R_4R_4'NC(=N(O)_nA_4)$ group (wherein R_4, R_4', n and A_4 are as defined above), an $A_4N=C(OR_2)$ group (wherein A_4 and R_2 are as defined above) or a NH_2CS group, and R_4 is as defined above,~~

~~—— a D_3 - R_4 group, wherein D_3 represents a nitro group or a R_4OSO_2 group (wherein R_4 is as defined above), and R_4 is as defined above, or~~

~~—— a R_4OSO_2 group, wherein R_4 is as defined above;~~


~~—— (2) the B group: an (a) group~~




~~in (a), E_1 and E_1' represent a methylene group optionally substituted with a C1-C10 alkyl group or a C1-C10 alkoxy group, or a carbonyl group, provided that E_1 and E_1' are not a carbonyl group at the same time, E_2 represents a C2-C10 alkylene group optionally substituted with an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a NR_4' group (wherein R_4' is as defined above), or a C3-C10 alkenylene group optionally substituted with an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a NR_4' group (wherein R_4' is as defined above), and R_4 is as defined above;~~

~~—— (3) the C group: a C2-C10 alkenyl group substituted with a halogen atom, a R_2B_4 group (wherein R_2 and B_4 are as defined above), a D_4R_4 group [wherein D_4 represents a hydroxyl group or an A_4O group (wherein A_4 is as defined above), and R_4 is as defined above], a D_5 group [wherein D_5 represents an $O=C(R_2)$ group (wherein R_2 is as defined above), an $A_4(O)_nN=C(R_2)$ group (wherein A_4, n and R_2 are as defined above), a $R_4B_0CO R_4(O)_nN=C(R_2)$ group (wherein R_4, R_4, n and R_2 are as defined above, and B_0 represents an oxy group, a thio group or a $N((O)_mR_4')$ group (wherein R_4' and m are as defined above)), a $D_2R_4(O)_nN=C(R_2)$ group (wherein D_2, R_4, n and R_2 are as defined above) or a $R_4A_4NN=C(R_2)$ group (wherein R_4, A_4 and R_2 are as defined above)], a $R_4A_4NO R_4$ group (wherein R_4, A_4 and R_4 are as defined above), a $R_4(A_4(O)_n)N$ group (wherein R_4, A_4 and n are as defined above), a D_2 group (wherein D_2 is as defined above) or a D_3 group (wherein D_3 is as defined above);~~

~~—— (4) the D group: a C2-C10 alkynyl group substituted with a (b) R_4 group [wherein, in (b)~~



~~G₁, G₂, G₄ and G₅ represent a methylene group which is connected with the adjacent atom via a single bond and which may be optionally substituted with a methyl group, or a methine group which is connected with the adjacent atom via a double bond and which may be optionally substituted with a methyl group, and G₃ represents a single bond, a double bond, a C1-C10 alkylene group optionally substituted with a methyl group, an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a NR₄ group (wherein R₄ is as defined above), or a C2-C10 alkenylene group optionally substituted with a methyl group, an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a NR₄ group (wherein R₄ is as defined above); and R₄ is as defined above], a (c) R₄ group (wherein, in (c)~~



~~J₁, J₂ and J₃ are the same or different, and represent a methine group optionally substituted with a methyl group, or a nitrogen atom; and R₄ is as defined above), a halogen atom, a R₂-B₄-R₄ group (wherein R₂, B₄ and R₄ are as defined above), a D₄-R₄ group (wherein D₄ and R₄ are as defined above), a D₅ group (wherein D₅ is as defined above), a D₁-R₄ group (wherein D₁ and R₄ are as defined above), a D₂ group (wherein D₂ is as defined above) or a D₃-R₄ group (wherein D₃ and R₄ are as defined above);~~

~~—— (5) the E group: an A₂-CO-R₅ group, provided that R₅ is not a vinylene group when A₂ is a hydroxyl group, wherein A₂ represents~~

~~—— (i) an A₃-B₄ group~~

~~—— wherein A₃ represents a hydrogen atom, or a C1-C10 alkyl group, or a C2-C10 haloalkyl group, or a C2-C10 alkenyl group optionally substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or R_a-(R₄)_m group (wherein R_a represents a phenyl group, a pyridyl group, a furyl group or a thienyl group, which may be optionally substituted with a halogen atom, a C1-C10 alkyl group, a C1-C10 alkoxy group or a nitro group, and R₄ and m are as defined above), or a C1-C10 alkyl group substituted with a~~
~~(b) R₄ group (wherein (b) and R₄ are as defined above), a (c) R₄ group (wherein (c) and R₄ are as defined above), a R₂-B₄-R₄ group (wherein R₂, B₄ and R₄ are as defined above), a D₄-R₄ group (wherein D₄ and R₄ are as defined above), a D₅ group (wherein D₅ is as defined above), a D₁-R₄ group (wherein D₁ and R₄ are as defined above), a D₂ group (wherein D₂ is as defined above), a D₃-R₄ group (wherein D₃ and R₄ are as defined above) or an A₄-SO₂-R₄ group {~~

~~wherein A_4 represents a (b) group (wherein (b) is as defined above), a (c) group (wherein (c) is as defined above) or a $R_4R_4'N$ group (wherein R_4 and R_4' are as defined above), and R_4 is as defined above}, and~~

~~—— B_4 represents an oxy group, a thio group or a $N((O)_mR_4)$ group (wherein R_4 and m are as defined above), provided that A_4 is not a hydrogen atom when B_4 is a thio group,~~

~~(ii) a $R_4B_4CO R_4B_4'$ group~~

~~—— wherein R_4 , B_4 and R_4' are as defined above, B_4' is the same as or different from B_4 and has the same meaning as B_4 has, provided that R_2 is not a hydrogen atom when B_4 is a thio group, or~~

~~—— a $D_2R_4B_4$ group, wherein D_2 , R_4 and B_4 are as defined above,~~

~~—— (iii) a $R_2SO_2NR_4$ group~~

~~—— wherein R_2 is as defined above, provided that a hydrogen atom is excluded; and R_4 is as defined above,~~

~~—— (iv) a (b) group, wherein (b) is as defined above,~~

~~—— (v) a (c) group, wherein (c) is as defined above, or~~

~~—— (vi) a $R_4A_4N NR_4'$ group, wherein R_4 , A_4 and R_4' are as defined above, and~~

~~R_5 represents a C2-C10 alkenylene group optionally substituted with a halogen atom, or a C2-C10 alkynylene group;~~

~~—— (6) the F group: an $A_5B_5R_6$ group~~

~~—— wherein A_5 represents a C2-C10 alkyl group substituted with a D_4 group (wherein D_4 is as defined above), a D_4 group (wherein D_4 is as defined above), a D_2 group (wherein D_2 is as defined above) or an A_4SO_2 group (wherein A_4 is as defined above), or a C1-C10 alkyl group substituted with a R_2B_4 group (wherein R_2 and B_4 are as defined above), a D_2 group (wherein D_2 is as defined above), a D_5 group (wherein D_5 is as defined above) or an A_2CO group (wherein A_2 is as defined above), B_5 represents a B_4 group (wherein B_4 is as defined above) or a NA_4 group (wherein A_4 is as defined above), and R_6 represents a single bond or a C1-C10 alkylene group;~~

~~—— (7) the G group: an $A_6B_5R_6$ group~~

~~wherein A_6 represents an (a) R_4 group (wherein (a) and R_4 are as defined above), or a C2-C10 alkenyl group, or a C2-C10 alkynyl group, or a C2-C10 alkenyl group substituted with a halogen atom, a R_2-B_1 group (wherein R_2 and B_1 are as defined above), a D_5 group (wherein D_5 is as defined above), a D_2 group (wherein D_2 is as defined above) or an A_2-CO group (wherein A_2 is as defined above), or a C2-C10 alkynyl group substituted with a halogen atom, a R_2-B_1 group (wherein R_2 and B_1 are as defined above), a D_5 group (wherein D_5 is as defined above), a D_2 group (wherein D_2 is as defined above) or an A_2-CO group (wherein A_2 is as defined above), or a C3-C10 alkenyl group substituted with a (b) group (wherein (b) is as defined above), a (c) group (wherein (c) is as defined above), a D_4 group (wherein D_4 is as defined above), a D_1 group (wherein D_1 is as defined above) or a D_3 group (wherein D_3 is as defined above), or a C3-C10 alkynyl group substituted with a D_4 group (wherein D_4 is as defined above), a D_1 group (wherein D_1 is as defined above) or a D_3 group (wherein D_3 is as defined above), and B_5 and R_6 are as defined above;~~

~~(8) the H group:~~

~~a $D_2-N((O)_n-A_1)-R_6$ group (wherein D_2 , n , A_1 and R_6 are as defined above),
a D_2 group (wherein D_2 is as defined above, provided that a cyano group is excluded),
a $R_4-(R_4'(O)_n)N-CR_4''-N-R_6$ group (wherein R_4 , R_4' , n and R_6 are as defined above, R_4'' is the same as or different from R_4 and has the same meaning as R_4 has),
a $R_4-(O)_n-N=CR_4'-NR_2-R_6$ group (wherein R_4 , n , R_4' , R_2 and R_6 are as defined above),
a $R_2-B_3-NR_4-CO-NR_4'-R_6$ group (wherein R_2 , B_3 , R_4 , R_4' and R_6 are as defined above),
a $D_2-CO-NR_4-R_6$ group (wherein D_2 , R_4 and R_6 are as defined above) or
an $A_2-COCO-NR_4-R_6$ group (wherein A_2 , R_4 and R_6 are as defined above);~~

~~(9) the I group:~~

~~an $A_7-B_6-N((O)_n-R_4)-R_6$ group [wherein A_7 represents a C2-C10 alkenyl group optionally substituted with a halogen atom, or a C2-C10 alkynyl group, or a C3-C10 haloalkynyl group, or a $R_2-B_1-R_4$ group (wherein R_2 , B_1 and R_4 are as defined above), or a D_4-R_4 group (wherein D_4 and R_4 are as defined above), or a D_5-R_4 group (wherein D_5 and R_4 are as defined above), or a D_1-R_4 group (wherein D_1 and R_4 are as defined above), or a (b) R_4 group (wherein (b) and R_4 are as defined above), or a (c) R_4 group (wherein (c) and R_4 are as defined above), or a D_2-R_4 group (wherein D_2 and R_4 are as defined above), or a D_3-R_4 group (wherein D_3 and R_4 are as defined above), or an $A_4-SO_2-R_4$ group (wherein A_4 and R_4 are as defined above), or an A_2-CO~~

~~R₄ group (wherein A₂ and R₄ are as defined above), B₆ represents a carbonyl group or a thiocarbonyl group, and n, R₁ and R₆ are as defined above],~~

~~an A₈ CS N((O)_nR₄) R₆ group [wherein A₈ represents a hydrogen atom or a C1-C10 alkyl group optionally substituted with a halogen atom, and n, R₁ and R₆ are as defined above],~~

~~— an A₂' B₂' B₂ N((O)_nR₄) R₆ group [wherein A₂' represents a C3-C10 alkenyl group optionally substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or a R₂-B₄-R₄' group (wherein R₂ and B₄ are as defined above, and R₄' represents a C2-C10 alkylene group), or a D₄-R₄' group (wherein D₄ and R₄' are as defined above), or a D₁-R₄' group (wherein D₁ and R₄' are as defined above), or a (b)-R₄' group (wherein (b) and R₄' are as defined above), or a (c)-R₄' group (wherein (c) and R₄' are as defined above), or a D₂-R₄ group (wherein D₂ and R₄ are as defined above), or a D₃-R₄' group (wherein D₃ and R₄' are as defined above), or an A₂ CO R₄ group (wherein A₂ and R₄ are as defined above), B₂' represents an oxy group, a thio group or a N((O)_nR₄') group (wherein n' is the same as or different from n and has the same meaning as n has, and R₄' is as defined above), and B₂, n, R₁ and R₆ are as defined above],~~

~~— an A₈' B₂' CS N((O)_nR₄) R₄ group [wherein A₈' represents a C1-C10 alkyl group or a C2-C10 haloalkyl group, B₂' is as defined above, and n, R₁ and R₆ are as defined above],~~

~~— an A₈' S B₂' N((O)_nR₄) R₆ group [wherein A₈', n, R₁ and R₆ are as defined above, and B₂' represents a carbonyl group or a sulfonyl group] or~~

~~— an A₂' SO₂ N((O)_nR₄) R₆ group [wherein A₂' represents a C2-C10 alkenyl group, or a C3-C10 alkenyl group substituted with a halogen atom, or a C3-C10 alkynyl group optionally substituted with a halogen atom, or a R₂-B₄-R₄' group (wherein R₂, B₄ and R₄' are as defined above), or a D₄-R₄' group (wherein D₄ and R₄' are as defined above), or a D₅-R₄ group (wherein D₅ and R₄ are as defined above), or a D₁-R₄' group (wherein D₁ and R₄' are as defined above), or a (b)-R₄' group (wherein (b) and R₄' are as defined above), or a (c)-R₄' group (wherein (c) and R₄' are as defined above), or a D₂-R₄ group (wherein D₂ and R₄ are as defined above), or a NO₂-R₄ group (wherein R₄ is as defined above), or an A₂ CO R₄ group (wherein A₂ and R₄ are as defined above), and n, R₁ and R₄ are as defined above];~~

~~— (10) the J group:~~

~~— an A₂ CO group (wherein A₂ is as defined above),~~

~~— an A₉ CS group (wherein A₉ represents A₂ or A₈),~~

~~an $A_9'(O)_mN=C(A_9)$ group (wherein A_9' represents A_1' or A_8' , and m and A_9 are as defined above);~~

~~— a D_2 -CO group (wherein D_2 is as defined above);~~

~~— an A_2 -COCO group (wherein A_2 is as defined above);~~

~~— an A_9 -CO- B_4' - R_6 group (wherein A_9 and R_6 are as defined above, and B_4' represents an oxy group or a thio group, provided that A_9 is not A_8 when B_4' is an oxy group);~~

~~— an A_9 -CS- B_4' - R_6 group (wherein A_9 , B_4' and R_6 are as defined above);~~

~~— an A_2'' -SO₂- B_4' - R_6 group (wherein A_2'' , B_4' and R_6 are as defined above);~~

~~— an A_8 -SO₂- B_4' - R_6 group (wherein A_8 , B_4' and R_6 are as defined above, provided that A_8 is not a hydrogen atom);~~

~~— an A_9' - B_2' - B_3 - B_4' - R_6 group (wherein A_9' , B_2' , B_3 , B_4' and R_6 are as defined above); or~~

~~— a C2-C10 alkenyl group substituted with a (b) group (wherein (b) is as defined above) or a (c) group (wherein (c) is as defined above);~~

~~— (11) the K group: an $A_{10}N((O)_nR_4)$ -CO- R_6 group~~

~~— wherein A_{10} represents a hydrogen atom (provided that n is not 0), an A_2'' -SO₂ group (wherein A_2'' is as defined above), an A_8 -SO₂ group (wherein A_8 is as defined above, provided that A_8 is not a hydrogen atom), an A_9' -O group (wherein A_9' is as defined above, provided that n is not 1), an A_9' group (wherein A_9' is as defined above, provided that A_8' is excluded when n is 0), a R_2 OCH₂ group (wherein R_2 is as defined above), an A_2 -CO- R_4 group (wherein A_2 and R_4 are as defined above) or an A_2 -CO-CH(CH₂CO- A_2) group (wherein A_2 is as defined above); and n, R_4 and R_6 are as defined above;~~

~~— (12) the L group:~~

~~— an $A_{10}'N((O)_nR_4)$ -SO₂- R_6 group [wherein A_{10}' represents a hydrogen atom (provided that n is not 0), an A_9' -O group (wherein A_9' is as defined above, provided that n is not 1), an A_9' group (wherein A_9' is as defined above, provided that A_8' is excluded when n is 0), a R_2 -CO group (wherein R_2 is as defined above), an A_2 -CO- R_4 group (wherein A_2 and R_4 are as defined above) or an A_2 -CO-CH(CH₂CO- A_2) group (wherein A_2 is as defined above); and n, R_4 and R_6 are as defined above];~~

~~— an $A_9''R_4N$ -SO₂- $N((O)_nR_4')$ - R_6 group [wherein A_9'' represents a hydrogen atom or an A_9' group (wherein A_9' is as defined above), and R_4 , n, R_4' and R_6 are as defined above] or~~

~~— a (b) $\text{SO}_2\text{N}((\text{O})_n\text{R}_4')$ R_6 -group [wherein (b), n, R_4' and R_6 are as defined above];~~
~~— (13) the M group:~~
~~— a $\text{R}_1(\text{R}_2\text{S})\text{C}=\text{N} \text{R}_6$ -group (wherein R_1 , R_2 and R_6 are as defined above);~~
~~— a $\text{R}_2\text{B}(\text{R}_2'\text{B}')\text{C}=\text{N} \text{R}_6$ -group (wherein R_2 and R_6 are as defined above, R_2' is the same as or different from R_2 and has the same meaning as R_2 has, and B and B' are the same or different and represent an oxy group or a thio group);~~
~~— a $\text{R}_1\text{R}_1'\text{N}(\text{R}_2\text{S})\text{C}=\text{N} \text{R}_6$ -group (wherein R_1 , R_1' , R_2 and R_6 are as defined above);~~
~~— a $\text{R}_1\text{N}=\text{C}(\text{SR}_2)\text{NR}_3' \text{R}_6$ -group (wherein R_1 , R_2 , R_3' and R_6 are as defined above) or~~
~~— a $\text{R}_1(\text{R}_1'\text{O})\text{N} \text{R}_6$ -group (wherein R_1 , R_1' and R_6 are as defined above);~~
~~— (14) the N group: an $\text{A}_{11}\text{P}(=\text{O})(\text{OR}_1') \text{R}_4$ -group~~
~~— wherein A_{11} represents a R_1 -group (wherein R_1 is as defined above), a $\text{R}_1\text{O} \text{R}_6$ -group (wherein R_1 and R_6 are as defined above) or a $\text{R}_1\text{OCO}-\text{CHR}_0$ -group (wherein R_1 and R_0 are as defined above), and R_1' and R_4 are as defined above;~~

~~III.(III)~~ in $(\text{Y}_A)_q$, Y_A is a substituent on a carbon atom and represents a group included in the following X group or Y group, q represents 0, 1, 2, 3 or 4, the sum of p (wherein p is as defined above) and q is 5 or less, Y_A s are the same or different when q is 2 or more, and the adjacent two same or different Y_A s together may form a group included in the Z group to be fused to the A ring when q is 2 or more,


(1) the X group: a M_a - group

wherein M_a represents a R_b - group (wherein R_b represents a C1-C10 alkyl group optionally substituted with a halogen atom), a halogen atom, a nitro group, a cyano group, a $\text{R}_c\text{-B}_a\text{-R}_d$ - group (wherein R_c represents a C1-C10 alkyl group optionally substituted with a halogen atom, B_a represents an oxy group, a thio group, a sulfinyl group or a sulfonyl group, and R_d represents a single bond or a C1-C10 alkylene group), a HOR_d - group (wherein R_d is as defined above), a $\text{R}_c\text{-CO-R}_d$ - group (wherein R_c represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom, and R_d is as defined above), a $\text{R}_c\text{-CO-O-R}_d$ - group (wherein R_c and R_d are as defined above), a $\text{R}_c\text{O-CO-R}_d$ - group (wherein R_c and R_d are as defined above), a HO-CO-CH=CH- group, a $\text{R}_c\text{R}_c'\text{N-R}_d$ - group (wherein R_c and R_c' are the same or different, R_c is as defined above, R_c' has the same meaning as R_c has, and R_d is as defined above), a $\text{R}_c\text{-CO-NR}_c'\text{-R}_d$ - group (wherein R_c , R_c' and R_d are as defined above), a $\text{R}_b\text{O-CO-}$


$N(R_e)-R_d$ - group (wherein R_b , R_e and R_d are as defined above), a $R_eR_e'N-CO-R_d$ - group (wherein R_e , R_e' and R_d are as defined above), a $R_eR_e'N-CO-NR_e''-R_d$ - group (wherein R_e , R_e' and R_e'' are the same or different, R_e and R_e' are as defined above, R_e'' has the same meaning as R_e has, and R_d is as defined above), a $R_eR_e'N-C(=NR_e'')-NR_e'''-R_d$ - group (wherein R_e , R_e' , R_e'' and R_e''' are the same or different, R_e , R_e' and R_e'' are as defined above, R_e''' has the same meaning as R_e has, and R_d is as defined above), a $R_b-SO_2-NR_e-R_d$ - group (wherein R_b , R_e and R_d are as defined above), a $R_eR_e'N-SO_2-R_d$ - group (wherein R_e , R_e' and R_d are as defined above), a C2-C10 alkenyl group or a C2-C10 alkynyl group;

(2) the Y group: a M_b-R_d -group, wherein M_b represents a M_c -group

~~wherein~~ wherein M_c represents a M_d-R_d' - group ~~wherein~~ wherein M_d represents a phenyl group optionally substituted with a M_a - group (wherein M_a is as defined above), a pyridyl group optionally substituted with a M_a - group (wherein M_a is as defined above), a naphthyl group optionally substituted with a M_a - group (wherein M_a is as defined above), a (b)- group (wherein (b) is as defined above), a (c)- group (wherein (c) is as defined above), a (d)- group



(wherein l is 2, 3 or 4, B_b represents an oxy group or a thio group) or an (e)- group



(wherein l and B_b are as defined above), and R_d' is the same as or different from R_d and has the same meaning as R_d ~~has~~ has),

a M_c-B_a - group (wherein M_c and B_a are as defined above), a M_c-CO - group (wherein M_c is as defined above), a M_c-CO-O - group (wherein M_c is as defined above), a M_cO-CO - group (wherein M_c is as defined above), a M_cR_eN - group (wherein M_c and R_e are as defined above), a $M_c-CO-NR_e$ - group (wherein M_c and R_e are as defined above), a $M_cO-CO-NR_e$ - group (wherein M_c and R_e are as defined above), a M_cR_eN-CO - group (wherein M_c and R_e are as defined above), a $M_cR_eN-CO-NR_e'$ - group (wherein M_c , R_e and R_e' are as defined above), a $M_cR_eN-C(=NR_e')$ - NR_e'' - group (wherein M_c , R_e , R_e' and R_e'' are as defined above), a $M_c-SO_2-NR_e$ - group (wherein M_c and R_e are as defined above) or a $M_cR_eN-SO_2$ - group (wherein M_c and R_e are as defined above), and

R_d is as defined above;

(3) the Z group:

a $-N=C(Y_a)-Y_a'$ - group (wherein Y_a represents a hydrogen atom, or a C1-C10 alkyl group optionally substituted with a halogen atom, or a C1-C10 alkoxy group, and Y_a' represents an oxy group, a thio group, or an imino group optionally substituted with a C1-C10 alkyl group),

a $-Y_b-Y_b'-Y_b''$ - group (wherein Y_b and Y_b'' are the same or different, and represent a methylene group, an oxy group, a thio group, a sulfinyl group, or an imino group optionally substituted with a C1-C10 alkyl group, and Y_b' represents a C1-C4 alkylene group optionally substituted with a halogen atom, or a C1-C4 alkylene group optionally having an oxo group) or

a $-Y_c-O-Y_c'-O$ - group (wherein Y_c and Y_c' are the same or different, and represent a C1-C10 alkylene group);


~~IV~~-(IV) Q_A represents a hydroxyl group, a (b)- group (wherein (b) is as defined above), an $A_9-B_6-B_c$ - group [~~wherein~~wherein A_9 and B_6 are as defined above, and B_c represents an oxy group or a $-N((O)_mR_1)$ - group (wherein m and R_1 are as defined above), provided that B_c is not a sulfonyl group when A_9 is a hydrogen atom], an $A_7''-SO_2-B_c$ - group (wherein A_7'' and B_c are as defined above), an $A_8-SO_2-B_c$ - group (wherein A_8 and B_c are as defined above, provided that A_8 is not a hydrogen atom), a $R_1R_1'-N-SO_2-B_c$ - group (wherein R_1 , R_1' and B_c are as defined above), a (b)- SO_2-B_c - group (wherein (b) and B_c are as defined above), an $A_9'-B_c$ - group (wherein A_9' and B_c are as defined above), a $D_5-R_4-B_c$ - group (wherein D_5 , R_4 and B_c are as defined above), a $M_c-B_3-B_c$ - group (wherein M_c , B_3 and B_c are as defined above) or a M_c-B_c - group (wherein M_c and B_c are as defined above);

~~V~~-(V) K_A represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, L_A represents a hydrogen atom, a C1-C10 alkyl group or a M_b -group (M_b is as defined above), or K_A and L_A may form a C1-C10 alkylene group or a $-C(M_a')=C(M_a'')-C(M_a''')=C(M_a'''')-$ group (M_a' , M_a'' , M_a''' and M_a'''' are the same or different, are the same as or different from M_a , and represent a hydrogen atom or M_a); and

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is

the same, selected substituents may be the same or different as long as they are selected within the range;range.

4. (Currently amended) A cinnamoyl compound represented by the formula (IV):



wherein:

A represents a benzene-phenyl ring or a pyridine-pyridyl ring,

X_a is a substituent on a carbon atom, and represents a_{3,1}-CH₂-CO-NH- (wherein a_{3,1} represents a C1-C10 alkoxy group) a C1-C10 alkyl group substituted with a cyano group; a C1-C10 alkyl group substituted with a tetrahydropyran 4-ylidene group; a C2-C10 alkenyl group substituted with a halogen atom or a cyano group; a C2-C10 alkenyl group substituted with a C1-C10 alkoxy carbonyl group; a C3-C10 alkynyl group substituted with a hydroxyl group; an a₀-r₁-b-r₁' group (wherein a₀ represents a methyl group substituted with a C1-C10 alkylthio group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r₂O-CO group (wherein r₂ represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxyl group), a carboxyl group, a rr'N-CO group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an a₁-NH-CO group (wherein a₁ represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an a₁'-CO group (wherein a₁' represents a morpholino group), a rr'N-CH₂ group (wherein r and r' are as defined above), a r₀(O)₁-CONH-CH₂ group (wherein r₀ represents a C1-C10 alkyl group, and l represents 0 or 1), a r-OCH₂ group (wherein r is as defined above), a r₀-CO group (wherein r₀ is as defined above), a cyano group, or a sulfomethyl group, r₁ represents a C1-C10 alkylene group, r₁' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a imino group}; an a₂-y-CO-NH group (wherein a₂ represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, and y represents an oxy group or an imino group); a r₀O-COCO-NH group (wherein r₀ is as defined above); an a₃-z-

~~NH- group (wherein a_2 represents a C2-C10 alkenyl group, or a C1-C10 alkyl group substituted with a C1-10 alkoxy group, a C1-C10 alkoxy carbonyl group, a carboxy group or a cyano group, and z represents a carbonyl group or a sulfonyl group); an a_4 -NHCO- group (wherein a_4 represents a C1-C10 alkoxy group, or a C3-C10 alkenyloxy group, or a r_0 -SO₂- group (wherein r_0 is as defined above), or a C2-C10 alkyl group substituted with a hydroxyl group or a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a r O-CO- group (wherein r is as defined above), a cyano group or an aminocarbonyl group, or a r O-CO-(r O-COCH₂)CH- group (wherein r is as defined above)); an a_5 -NHSO₂- group (wherein a_5 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group); a r_0 ON-CH- group (wherein r_0 is as defined above); a r_0 NHCSNH- group (wherein r_0 is as defined above); a r_0 NHC(Sr_0')=N- group (wherein r_0 is as defined above, r_0' is the same as the different from r_0 and has the same meaning as r_0 has); or a (r_0 O)₂P(=O)CH₂- group (wherein r_0 is as defined above);~~

p represents 1, 2 or 3, and when p is 2 or more, X_as are the same or different;

Y_a represents a halogen atom, a nitro group, a r_0 CO-NH- group (wherein r_0 is a C1-C10 alkyl group as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group;

q represents 0, 1 or 2, and when q is 2 or more, Y_as are the same or different;


q_a represents a r_a -O- group {wherein r_a represents a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with a r_0r_0' N-CH₂- group (wherein r_0 ~~and r_0' are~~ is as defined above and r_0' is the same as the different from r_0 and has the same meaning as r_0 has), a r OCH₂- group (wherein r is a hydrogen atom or a C1-C10 alkyl group as defined above), a r_0 -CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, a carboxy group, an aminocarbonyl group or a cyano group, or a r_3 - r_1 - group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 is a C1-C10 alkylene group as defined above)}; a piperidino group; a morpholino group; or a r_4r_4' N- group (wherein r_4 and r_4' are the same or different, and represent a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that r_4 and r_4' are not a hydrogen atom at the same time);

K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group; or

K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group;

the term “as defined above” used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range;range.

5. (Currently amended) A cinnamoyl compound represented by the formula (V):



wherein:

a represents a benzene-phenyl ring or a pyridine-pyridyl ring;

x represents a methine group or a nitrogen atom;

X_a is a substituent on a carbon atom, and represents a₃₋₁-CH₂-CO-NH- (wherein a₃₋₁ represents a C1-C10 alkoxy group) ~~a C1-C10 alkyl group substituted with a cyano group; a C1-C10 alkyl group substituted with a tetrahydropyran-4-ylidene group; a C2-C10 alkenyl group substituted with a halogen atom or a cyano group; a C2-C10 alkenyl group substituted with a C1-C10 alkoxy carbonyl group; a C3-C10 alkynyl group substituted with a hydroxyl group; an a₀-r₁-b-r₄' group (wherein a₀ represents a methyl group substituted with a C1-C10 alkylthio group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r₃O-CO group (wherein r₂ represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxyl group), a carboxyl group, a rr'N-CO group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an a₄-NH-CO group (wherein a₄ represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an a₄'-CO group (wherein a₄' represents a morpholino group), a rr'N-CH₂ group (wherein r and r' are as defined above), a r₀(O)-CONH-CH₂ group (wherein r₀ represents a C1-C10 alkyl group, and l represents 0 or 1), a r-OCH₂ group (wherein r is as defined above), a r₀-CO group (wherein r₀ is~~

as defined above), a cyano group, or a sulfomethyl group, r_1 represents a C1-C10 alkylene group, r_1' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio group, a sulfinyl group, a sulfonyl group or an imino group}; an a_2 -y-CO-NH- group (wherein a_2 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, and y represents an oxy group or an imino group); a r_0 O-COCO-NH- group (wherein r_0 is as defined above); an a_3 -z-NH- group (wherein a_3 represents a C2-C10 alkenyl group, or a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, a C1-C10 alkoxycarbonyl group, a carboxy group or a cyano group, and z represents a carbonyl group or a sulfonyl group); an a_4 -NHCO- group (wherein a_4 represents a C1-C10 alkoxy group, or a C3-C10 alkenyloxy group, or a r_0 -SO₂- group (wherein r_0 is as defined above), or a C2-C10 alkyl group substituted with a hydroxyl group or a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a r O-CO- group (wherein r is as defined above), a cyano group or an aminocarbonyl group, or a r O-CO-(r O-COCH₂)CH- group (wherein r is as defined above)); an a_5 -NHSO₂- group (wherein a_5 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group); a r_0 ON=CH- group (wherein r_0 is as defined above); a r_0 NHCSNH- group (wherein r_0 is as defined above); a r_0 NHC(-S r_0')=N- group (wherein r_0 is as defined above, r_0' is the same as the different from r_0 and has the same meaning as r_0 has); or a (r_0 O)₂P(=O)CH₂- group (wherein r_0 is as defined above);

p represents 1, 2 or 3, and when p is 2 or more, X_as are the same or different;

Y_a represents a halogen atom, a nitro group, a r_0 CO-NH- group (wherein r_0 is a C1-C10 alkyl group as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group;

q represents 0, 1 or 2, and when q is 2 or more, Y_as are the same or different;

q_a represents a r_a -O- group {wherein r_a represents a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with a r_0r_0' N-CH₂- group (wherein r_0 and r_0' are as defined above and r_0' is the same as the different from r_0 and has the same meaning as r_0 has), a r OCH₂- group (wherein r is a hydrogen atom or a C1-C10 alkylene group as defined above), a r_0 -CO- group (wherein r_0 is as defined above), a C1-C10 alkoxycarbonyl group, a carboxy group, an aminocarbonyl group or a cyano group, or a r_3 - r_1 -group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 is a C1-C10 alkylene group as defined above)}; a piperidino group; a morpholino group; or a r_4r_4' N- group (wherein r_4 and r_4' are the same or different, and represent a hydrogen atom, a C1-C10 alkyl group, a C3-

C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that r_4 and r_4' are not a hydrogen atom at the same time);


t_a represents a r_b - group (wherein r_b is the same as or different from r_a , and has the same meaning as r_a has) or a r_3' - group (wherein r_3' is the same as or different from r_3 , and has the same meaning as r_3 has);

K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group; or

K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group;

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range;range.

6. (Currently amended) A 2H-pyran-2-one compound represented by the formula (VI):



wherein:

a represents a ~~benzene-phenyl~~ ring or a ~~pyridine-pyridyl~~ ring;

x represents a methine group or a nitrogen atom;

X_a is a substituent on a carbon atom, and represents a_{3-1} -CH₂-CO-NH- (wherein a_{3-1} ~~represents a C1-C10 alkoxy group; a C1-C10 alkyl group substituted with a cyano group; a C1-C10 alkyl group substituted with a tetrahydropyran-4-ylidene group; a C2-C10 alkenyl group substituted with a halogen atom or a cyano group; a C2-C10 alkenyl group substituted with a C1-C10 alkoxy carbonyl group; a C3-C10 alkynyl group substituted with a hydroxyl group; an a_0 - r_4 - b - r_4' group (wherein a_0 represents a methyl group substituted with a C1-C10 alkylthio group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a~~

~~C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r_2 O-CO group (wherein r_2 represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxyl group), a carboxyl group, a rr' N-CO group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an a_1 -NH-CO group (wherein a_1 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an a_1' -CO group (wherein a_1' represents a morpholino group), a rr' N-CH₂ group (wherein r and r' are as defined above), a r_0 (O)₁-CONH-CH₂ group (wherein r_0 represents a C1-C10 alkyl group, and 1 represents 0 or 1), a r -OCH₂ group (wherein r is as defined above), a r_0 -CO group (wherein r_0 is as defined above), a cyano group, or a sulfomethyl group, r_1 represents a C1-C10 alkylene group, r_1' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a imino group}; an a_2 -y-CO-NH group (wherein a_2 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, and y represents an oxy group or an imino group); a r_0 O-COCO-NH group (wherein r_0 is as defined above); an a_3 -z-NH group (wherein a_3 represents a C2-C10 alkenyl group, or a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, a C1-C10 alkoxy carbonyl group, a carboxy group or a cyano group, and z represents a carbonyl group or a sulfonyl group); an a_4 -NHCO group (wherein a_4 represents a C1-C10 alkoxy group, or a C3-C10 alkenyloxy group, or a r_0 -SO₂ group (wherein r_0 is as defined above), or a C2-C10 alkyl group substituted with a hydroxyl group or a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a r O-CO group (wherein r is as defined above), a cyano group or an aminocarbonyl group, or a r O-CO-(r O-COCH₂)CH group (wherein r is as defined above)); an a_5 -NHSO₂ group (wherein a_5 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group); a r_0 ON=CH group (wherein r_0 is as defined above); a r_0 NHCSNH group (wherein r_0 is as defined above); a r_0 NHC(Sr_0')-N group (wherein r_0 is as defined above, r_0' is the same as the different from r_0 and has the same meaning as r_0 has); or a (r_0 O)₂P(=O)CH₂ group (wherein r_0 is as defined above);~~

p represents 1, 2 or 3, and when p is 2 or more, X_a s are the same or different;

Y_a represents a halogen atom, a nitro group, a r_0 CO-NH- group (wherein r_0 is a C1-C10 alkyl group as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group;


q represents 0, 1 or 2, and when q is 2 or more, Y_a s are the same or different;

q_a represents a r_a -O- group {wherein r_a represents a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with

a r_0r_0' -N-CH₂- group (wherein r_0 and r_0' are as defined above and r_0' is the same as the different from r_0 and has the same meaning as r_0 has), a r OCH₂- group (wherein r is a hydrogen atom or a C1-C10 alkyl group as defined above), a r_0 -CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, a carboxy group, an aminocarbonyl group or a cyano group, or a r_3 - r_1 - group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 is a C1-C10 alkylene group as defined above)); a piperidino group; a morpholino group; or a r_4r_4' -N- group (wherein r_4 and r_4' are the same or different, and represent a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that r_4 and r_4' are not a hydrogen atom at the same time);

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range.

7. (Currently amended) A 2H-pyran-2-one compound represented by the formula (VII):




wherein:

X_a' represents $a_{3,1}$ -CH₂-CO-NH- (wherein $a_{3,1}$ represents a C1-C10 alkoxy group) a C1-C10 alkyl group substituted with a cyano group, or a C2-C10 alkenyl group substituted with a halogen atom or a cyano group, or an a_0' - r_1 -O group (a_0' represents a methyl group substituted with a C1-C10 alkylthio group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a HOCH₂- group or a cyano group, and r_1 represents a C1-C10 alkylene group), or an a_6 -CONH- group (a_6 represents a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, or a C2-C10 alkoxy group substituted with a C1-C10 alkoxy group), or an a_7 -NHCO- group (a_7 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a

~~C1-C10 alkoxy carbonyl group); q_a' represents an amino group substituted with a C3-C10 alkynyl group, a piperidino group, a morpholino group or a r_a' -O group (r_a' represents a hydrogen atom, a C1-C10 alkyl group or a C3-C10 alkenyl group).~~

8. (Currently amended) A 2H-1-benzopyran-2-one compound represented by the formula (VIII):



wherein:

a represents a ~~benzene-phenyl~~ ring or a ~~pyridine-pyridyl~~ ring;

x represents a methine group or a nitrogen atom;

X_a is a substituent on a carbon atom, and represents ~~$a_{3,1}$ -CH₂-CO-NH-~~ (wherein $a_{3,1}$ represents a C1-C10 alkoxy group) ~~a C1-C10 alkyl group substituted with a cyano group; a C1-C10 alkyl group substituted with a tetrahydropyran-4-ylidene group; a C2-C10 alkenyl group substituted with a halogen atom or a cyano group; a C2-C10 alkenyl group substituted with a C1-C10 alkoxy carbonyl group; a C3-C10 alkynyl group substituted with a hydroxyl group; an a_0 - r_1 - b - r_1' group (wherein a_0 represents a methyl group substituted with a C1-C10 alkylthio group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r_2 O-CO group (wherein r_2 represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxyl group), a carboxyl group, a rr' N-CO group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an a_1 -NH-CO group (wherein a_1 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an a_1' -CO group (wherein a_1' represents a morpholino group), a rr' N-CH₂ group (wherein r and r' are as defined above), a r_0 (O)-CONH-CH₂ group (wherein r_0 represents a C1-C10 alkyl group, and l represents 0 or 1), a r OCH₂ group (wherein r is as defined above), a r_0 -CO group (wherein r_0 is as defined above), a cyano group, or a sulfomethyl group, r_1 represents a C1-C10 alkylene group, r_1' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio~~

~~group, a sulfinyl group, a sulfonyl group or an imino group}; an a_2 -y CO-NH group (wherein a_2 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, and y represents an oxy group or an imino group); a r_0 O-COCO-NH group (wherein r_0 is as defined above); an a_3 -z-NH group (wherein a_3 represents a C2-C10 alkenyl group, or a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, a C1-C10 alkoxy carbonyl group, a carboxy group or a cyano group, and z represents a carbonyl group or a sulfonyl group); an a_4 -NHCO group (wherein a_4 represents a C1-C10 alkoxy group, or a C3-C10 alkenyloxy group, or a r_0 -SO₂ group (wherein r_0 is as defined above), or a C2-C10 alkyl group substituted with a hydroxyl group or a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a r O-CO group (wherein r is as defined above), a cyano group or an aminocarbonyl group, or a r O-CO (r O-COCH₂)CH- group (wherein r is as defined above)); an a_5 -NHSO₂ group (wherein a_5 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group); a r_0 ON=CH group (wherein r_0 is as defined above); a r_0 NHCSNH group (wherein r_0 is as defined above); a r_0 NHC(Sr_0')=N group (wherein r_0 is as defined above, r_0' is the same as the different from r_0 and has the same meaning as r_0 has); or a $(r_0O)_2P(=O)CH_2$ group (wherein r_0 is as defined above);~~

p represents 1, 2 or 3, and when p is 2 or more, X_a s are the same or different;

Y_a represents a halogen atom, a nitro group, a r_0 CO-NH- group (wherein r_0 is a C1-C10 alkyl group as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group;

q represents 0, 1 or 2, and when q is 2 or more, Y_a s are the same or different;

q_a represents a r_a -O- group {wherein r_a represents a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with a r_0r_0' N-CH₂- group (wherein r_0 and r_0' are is as defined above and r_0' is the same as the different from r_0 and has the same meaning as r_0 has), a r OCH₂- group (wherein r is a hydrogen atom or a C1-C10 alkyl group as defined above), a r_0 -CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, a carboxy group, an aminocarbonyl group or a cyano group, or a r_3 - r_1 - group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 is a C1-C10 alkylene group as defined above)}; a piperidino group; a morpholino group; or a r_4r_4' N- group (wherein r_4 and r_4' are the same or different, and represent a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that r_4 and r_4' are not a hydrogen atom at the same time);

t_a represents a r_b - group (wherein r_b is the same as or different from r_a , and has the same meaning as r_a has) or a r_3' - group (wherein r_3' is the same as or different from r_3 , and has the same meaning as r_3 has);


K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group; or

K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group;

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range;range.

9. (Currently amended) A 2H-1-benzopyran-2-one compound represented by the formula (IX):


(IX)



wherein:

X_a'' represents ~~a_{3-1} -CH₂-CO-NH-~~ (wherein ~~a_{3-1}~~ represents a C1-C10 alkoxy group) ~~a C1-C10 alkoxy group substituted with a cyano group or a hydroxymethyl group, or an a_6 -CONH group (a_6 represents a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, or a C2-C10 alkoxy group substituted with a C1-C10 alkoxy group), or an a_7 -NHCO group (a_7 represents a C2-C10 alkyl group substituted with a hydroxy group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a C1-C10 alkoxy carbonyl group), and q_a'' represents a hydroxy group, a C1-C10 alkoxy group or a piperidino~~ group;group.


10. (Currently amended) A 2H-pyran-2-one compound represented by the formula (X):



wherein:

~~X₁ represents a₃₋₁-CH₂-CO-NH- (wherein a₃₋₁ represents a C1-C10 alkoxy group) a C2-C4 alkenyl group substituted with a cyano group, an A₁-R₁-O group (A₁ represents a C1-C4 alkylthio group, a C2-C4 alkenyl group, a C2-C4 alkynyl group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group, and R₁ represents a C1-C4 alkylene group), an A_{II}-(y)_m-z-NH group (A_{II} represents a C2-C4 alkenyl group, or a C1-C4 alkyl group substituted with a C1-C4 alkoxy group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group, y represents an oxy group or an imino group, z represents a carbonyl group or a sulfonyl group, and m represents 0 or 1) or an A_{III}-NHCO group (A_{III} represents a methanesulfonyl group, or a C1-C4 alkyl group substituted with a hydroxy group, a C1-C4 alkoxy group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group), a_I represents a hydroxy group, a C1-C4 alkoxy group, a C2-C4 alkenyloxy group, a C2-C4 alkynyloxy group, a C1-C4 alkylamino group, a C2-C4 alkenylamino group, a C2-C4 alkynylamino group, a morpholino group or a piperidino group, Y_I represents a halogen atom, a nitro group, a C1-C4 alkyl group or a C1-C4 alkoxy group, n represents 0, 1 or 2 and, when n is 2, Y_Is may be different; different.~~


11. (Currently amended) A 2H-1-benzopyran-2-one compound represented by the formula (XI):



wherein:

X_I represents $a_{3-I}-CH_2-CO-NH-$ (wherein a_{3-I} represents a C1-C10 alkoxy group) ~~a C2-C4 alkenyl group substituted with a cyano group, an A_I-R_I-O group (A_I represents a C1-C4 alkylthio group, a C2-C4 alkenyl group, a C2-C4 alkynyl group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group, and R_I represents a C1-C4 alkylene group), an $A_{II}(y)_m-z-NH$ group (A_{II} represents a C2-C4 alkenyl group, or a C1-C4 alkyl group substituted with a C1-C4 alkoxy group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group, y represents an oxy group or an imino group, z represents a carbonyl group or a sulfonyl group, and m represents 0 or 1) or an $A_{III}-NHCO$ group (A_{III} represents a methanesulfonyl group, or a C1-C4 alkyl group substituted with a hydroxy group, a C1-C4 alkoxy group, a C1-C4 alkoxy carbonyl group, a carboxy group or a cyano group), a_I represents a hydroxy group, a C1-C4 alkoxy group, a C2-C4 alkenyloxy group, a C2-C4 alkynyloxy group, a C1-C4 alkylamino group, a C2-C4 alkenylamino group, a C2-C4 alkynylamino group, a morpholino group or a piperidino group, Y_I represents a halogen atom, a nitro group, a C1-C4 alkyl group or a C1-C4 alkoxy group, n represented 0, 1 or 2 and, when n is 2, Y_I 's Y_I 's may be different; different.~~


12. (Currently amended) A 2H-pyran-2-one compound represented by the formula (XII):



wherein:

X_{II} represents $a_{3-II}-CH_2-CO-NH-$ (wherein a_{3-II} represents a C1-C10 alkoxy group) ~~an allyloxy group, a propargyloxy group, a cyanomethoxy group, a methoxyacetyl amino group, a methoxycarbonylmethylaminocarbonyl group or a 2-cyanoethenyl group, and a_{II} represents a hydroxy group, a methoxy group or a morpholino group; group.~~

13. (Currently amended) A 2H-1-benzopyran-2-one compound represented by the formula (XIII):




wherein:

X_{II}' represents $a_{3-1}-CH_2-CO-NH-$ (wherein a_{3-1} represents a C1-C10 alkoxy group), a cyanomethoxy group, a methoxyacetyl amino group or a 2-hydroxyethylaminocarbonyl group;


14-17. (Cancelled).

18. (Currently amended) The 2H-pyran-2-one compound according to claim 1 represented by the formula (XVIII):



19-24. (Cancelled)


25. (Currently amended) The 2H-1-benzopyran-2-one compound according to claim 1 represented by the formula (XXV):



26-46. (Cancelled)


47. (Withdrawn-currently amended) A process for producing a cinnamoyl compound represented by the formula (XLVII''):

(XLVII'')



wherein A, X_c, Y_a, p, q, r_c, K_a and L_a are as defined below, and the term “as defined above (or below)” used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above (or below) and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range; which comprises reacting a cinnamoyl compound represented by the formula (XLVII):

(XLVII)



wherein

A represents a ~~benzene-phenyl~~ ring or a ~~pyridine-pyridyl~~ ring,

X_c is a substituent on a carbon atom, and represents ~~a₃₋₁-CH₂-CO-NH-~~ (wherein ~~a₃₋₁~~ represents a C1-C10 alkoxy group) ~~a C1-C10 alkyl group substituted with a cyano group; a C1-C10 alkyl group substituted with a tetrahydropyran-4-ylidene group; a C2-C10 alkenyl group substituted with a halogen atom or a cyano group; a C2-C10 alkenyl group substituted with a C1-C10 alkoxy carbonyl group; a C2-C10 alkynyl group substituted with a hydroxymethyl group; an a₀₋₁-F₁-b-F₁' group (wherein a₀₋₁ represents a methyl group substituted with a C1-C10 alkylthio~~

~~group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r_2 O-CO group (wherein r_2 represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxyl group), a rr' N-CO group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an a_4 -NH-CO group (wherein a_4 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an a_4' -CO group (wherein a_4' represents a morpholino group), a rr' N-CH₂ group (wherein r and r' are as defined above), a r_0 (O)_l-CONH-CH₂ group (wherein r_0 represents a C1-C10 alkyl group, and l represents 0 or 1), a r OCH₂ group (wherein r is as defined above), a r_0 -CO group (wherein r_0 is as defined above), or a cyano group, r_1 represents a C1-C10 alkylene group, r_1' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio group, a sulfinyl group, a sulfonyl group or a imino group}; an a_2 -y-CO-NH group (wherein a_2 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, and y represents an oxy group or an imino group); a r_0 O-COCO-NH group (wherein r_0 is as defined above); an a_2 -z-NH group (wherein a_2 represents a C2-C10 alkenyl group, or a C1-C10 alkyl group substituted with a C1-C10 alkoxy group, a C1-C10 alkoxy carbonyl group or a cyano group, and z represents a carbonyl group or a sulfonyl group); an a_4 -NHCO group (wherein a_4 represents a C1-C10 alkoxy group, or a C3-C10 alkenyloxy group, or a r_0 -SO₂ group (wherein r_0 is as defined above), or a C2-C10 alkyl group substituted with a hydroxyl group or a C1-C10 alkoxy group, or a C1-C10 alkyl group substituted with a r_0 O-CO group (wherein r_0 is as defined above), a cyano group or an aminocarbonyl group, or a r_0 O-CO (r_0 O-COCH₂)CH group (wherein r_0 is as defined above)); an a_5 -NH-SO₂ group (wherein a_5 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group); a r_0 ON=CH group (wherein r_0 is as defined above); a r_0 NHCSNH group (wherein r_0 is as defined above); a r_0 NHC(Sr_0')=N group (wherein r_0 is as defined above, r_0' is the same as the different from r_0 and has the same meaning as r_0 has); or a (r_0 O)₂P(=O)CH₂ group (wherein r_0 is as defined above);~~

p represents 1, 2 or 3, and when p is 2 or more, X_c s are the same or different;

Y_a represents a halogen atom, a nitro group, a r_0 CO-NH- group (wherein r_0 is a C1-C10 alkyl group as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group;

q represents 0, 1 or 2, and when q is 2 or more, Y_a s are the same or different;

K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group, or

K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group, and


the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range, with a compound represented by the formula (XLVII'):

r_c -V (XLVII')

wherein r_c represents a t_c '-group {wherein t_c ' represents a C1-C10 alkyl group; a C3-C10 alkenyl group; a C3-C10 alkynyl group; a C1-C10 alkyl group substituted with a r_0r_0' N-CH₂- group (wherein r_0 and r_0' are as defined above and r_0' is the same as the different from r_0 and has the same meaning as r_0 has), a $rOCH_2$ - group (wherein r is as defined above a hydrogen atom or a C1-C10 alkyl group), a r_0 -CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, an aminocarbonyl group or a cyano group; or a r_3 - r_1 - group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 is a C1-C10 alkylene group as defined above)}, and V represents a leaving group, and

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range; range.

48. (Withdrawn-currently amended) A process for producing a cinnamoyl compound represented by the formula (XLVIII'):



wherein:

A is as defined below,

X_d' is a substituent on a carbon atom, and represents a_{3-1} -CH₂-CO-NH- (wherein a_{3-1} represents a C1-C10 alkoxy group) ~~an a_{0d} -r₁-b-r₁' group (wherein a_{0d} ' represents a carboxy group, and r₁, r₁' and b are as defined below), a HO-COCO-NH group, an a_{1d} -z-NH group (wherein a_{1d} ' represents a C1-C10 alkyl group substituted with a carboxy group, and z is as defined below), or an a_{1d} -NHCO group (wherein a_{1d} ' represents a C1-C10 alkyl group substituted with a carboxy group, or a HO-CO-(HO-COCH₂)CH group),~~

p is as defined below and, and when p is 2 or more, X_d 's are the same or different,

Y_a and q are as defined below,


q_d' represents a r_d'' -O- group {wherein r_d'' represents a hydrogen atom; a C1-C10 alkyl group; a C3-C10 alkenyl group; a C3-C10 alkynyl group; a C1-C10 alkyl group substituted with a r_0r_0' -N-CH₂- group (wherein r_0 and r_0' are as defined below), a $rOCH_2$ - group (wherein r is as defined below), a r_0 -CO- group (wherein r_0 is as defined below), a carboxy group, an aminocarbonyl group or a cyano group; or a r_3 -r₁- group (wherein r_3 represents a phenyl group or a pyridyl group, and r₁ is as defined below)}, a piperidino group, a morpholino group, or a r_4r_4' -N- group (wherein r_4 and r_4' are as defined below, provided that they are not hydrogen atom at the same time),

K_a and L_a are as defined below, and

the term "as defined above (or below)" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above (or below) and, among the plural substituents, although the selection range of

substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range;

which comprises hydrolyzing a cinnamoyl compound represented by the formula (XLVIII):



wherein:

A represents a benzene ring or a pyridine ring,

X_d is a substituent on a carbon atom, and represents an $a_{0d}-r_1-b-r_1'$ - group {wherein a_{0d} represents a r_2O-CO- group (wherein r_2 represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxy group), r_1 represents a C1-C10 alkylene group, r_1' represents a single bond or a C1-C10 alkylene group, and b represents an oxy group, a thio group, a sulfinyl group, a sulfonyl group or an imino group}, a $r_0O-COCO-NH-$ group (wherein r_0 represents a C1-C10 alkyl group), an $a_{3d}-z-NH-$ group (wherein a_{3d} represents a C1-C10 alkyl group substituted with a C1-C10 alkoxy carbonyl group, and z represents a carbonyl group or a sulfonyl group), or an $a_{4d}-NHCO-$ group {wherein a_{4d} represents a C1-C10 alkyl group substituted with a r_0O-CO- group (wherein r_0 is as defined above), or a $r_0O-CO-(r_0O-COCH_2)CH-$ group (wherein r_0 is as defined above)},

p represents 1, 2 or 3, and when p is 2 or more, X_{ds} are the same or different,

Y_a represents a halogen atom, a nitro group, a $r_0CO-NH-$ group (wherein r_0 is as defined above), a C1-C10 alkyl group or a C1-C10 alkoxy group,

q represents 0, 1 or 2, and when q is 2 or more, Y_{as} are the same or different;


Q_d represents a r_d-O- group {wherein r_d represents a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with a $r_0r_0'N-CH_2-$ group (wherein r_0 is as defined above, and r_0' is the same as or different from r_0 and has the same meaning as r_0 has), a $rOCH_2-$ group (wherein r is as defined above), a r_0-CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, a carboxy group, an aminocarbonyl group or a cyano group, or a r_3-r_1 -group (wherein r_3 represents a phenyl group or

a pyridyl group, and r_1 is as defined above)); a piperidino group; a morpholino group; or a r_4r_4' N- group (wherein r_4 and r_4' represent a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that they are not a hydrogen atom at the same time),.

K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group, or


K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group, the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the ~~range;range~~.

49. (Withdrawn-currently amended) A process for producing a cinnamoyl compound represented by the formula (XLIX'')



wherein ~~X_e' represents~~ is a substituent on a carbon atom and represents ~~a_{3-1} -CH₂-CO-NH-~~ (wherein ~~a_{3-1} represents a C1-C10 alkoxy group~~)~~an a_{0e}' - r_1 - b group {wherein a_{0e}' represents an a_{0e} group (wherein a_{0e} is as defined below), a 3-sulfopropyl group or a 4-sulfobutyl group, and r_1 and b are as defined below}~~, and A, Y_a , p, q, q_e , K_a and L_a are as defined below, and the term "as defined above (or below)" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above (or below) and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the range;

which comprises reacting a cinnamoyl compound represented by the formula (XLIX):



wherein:

A represents a benzene ring or a pyridine ring,

X_e is a substituent on a carbon atom, and represents a H- b'' - group (wherein b'' represents an oxy group or a thio group),

p represents 1, 2 or 3 and, when p is 2 or more, X_e s are the same or different,

Y_a represents a halogen atom, a nitro group, a $r_0CO-NH-$ group (wherein r_0 is a C1-C10 alkyl group), a C1-C10 alkyl group or a C1-C10 alkoxy group,

q represents 0, 1 or 2, and when q is 2 or more, Y_a s are the same or different;

Q_e represents a r_e-O- group {wherein r_e represents a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, a C1-C10 alkyl group substituted with a $r_0r_0'N-CH_2-$ group (wherein r_0 is as defined above, and r_0' is the same as or different from r_0 and has the same meaning as r_0 has), a $rOCH_2-$ group (wherein r represents a hydrogen atom or a C1-C10 alkyl group), a r_0-CO- group (wherein r_0 is as defined above), a C1-C10 alkoxy carbonyl group, an aminocarbonyl group or a cyano group, or a r_3-r_1 -group (wherein r_3 represents a phenyl group or a pyridyl group, and r_1 represents a C1-C10 alkylene group)}; a piperidino group; a morpholino group; or a $r_4r_4'N-$ group (wherein r_4 and r_4' represent a hydrogen atom, a C1-C10 alkyl group, a C3-C10 alkenyl group, a C3-C10 alkynyl group, or a C2-C10 alkyl group substituted with a C1-C10 alkoxy group, provided that they are not a hydrogen atom at the same time),

K_a represents a hydrogen atom, a halogen atom or a C1-C10 alkyl group, and L_a represents a hydrogen atom or a C1-C10 alkyl group, or

K_a and L_a together may form a C1-C10 alkylene group or a 1,3-butadienylene group, and

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is

the same, selected substituents may be the same or different as long as they are selected within the range,

with a compound represented by the formula (XLIX'):

$a_0e-r_1''-V'$ (XLIX')

wherein

a_0e represents a methyl group substituted with a C1-C10 alkylthio group, a methyl group substituted with a C1-C10 alkylsulfinyl group, a methyl group substituted with a C1-C10 alkylsulfonyl group, a C2-C10 alkenyl group, a C2-C10 alkynyl group, a r_2O-CO- group (wherein r_2 represents a C1-C10 alkyl group, or a C2-C10 alkyl group substituted with a hydroxy group), a $rr'N-CO-$ group (wherein r and r' are the same or different, and represent a hydrogen atom or a C1-C10 alkyl group), an $a_1-NH-CO-$ group (wherein a_1 represents a C2-C10 alkyl group substituted with a C1-C10 alkoxy group), an $a_1'-CO-$ group (wherein a_1' represents a morpholino group), a $rr'N-CH_2-$ group (wherein r is as defined above, r' is the same as or different from r and has the same meaning as r has), a $r_0-(O)_l-CONH-CH_2-$ group (wherein r_0 is as defined above, and l represents 0 or 1), a $r-OCH_2-$ group (wherein r is as defined above), a r_0-CO- group (wherein r_0 is as defined above) or a cyano group,

r_1'' is the same as or different from r_1 and has the same meaning as r_1 has, and V' represents a leaving group or a hydroxy group, or 1,3-propanesultone or 1,4-butanedisultone

the term "as defined above" used for the same symbols among plural substituents means that the plural substituents independently represent the same meaning as that described above and, among the plural substituents, although the selection range of substituents to be selected is the same, selected substituents may be the same or different as long as they are selected within the ~~range;~~range.

50. (Cancelled)

51. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 1 and an inert ~~carrier;~~carrier.

52. (Cancelled)

53. (Withdrawn-currently amended) A composition for improving tissue fibrosis, which comprises a compound according to claim 1 and an inert ~~carrier~~carrier.

54. (Withdrawn-currently amended) A method for improving tissue fibrosis, which comprises administering an effective amount of a compound according to claim 1 to a mammal in need ~~thereof~~thereof.

55. (Cancelled)

56. (Withdrawn-currently amended) A composition for suppressing the activity of TGF- β , which comprises a compound according to claim 1 and an inert ~~carrier~~carrier.

57. (Cancelled)

58. (Withdrawn-currently amended) A composition for hair growth which comprises a compound according to claim 1 and an inert ~~carrier~~carrier.

59. (Withdrawn-currently amended) A method for growing hair, which comprises administering an effective amount of a compound according to claim 1 to a mammal in need ~~thereof~~thereof.

60. (Cancelled)

61. (Withdrawn-currently amended) An agent for treating chronic renal failure, which comprises a compound according to claim 1 and an inert ~~carrier~~carrier.

62. (Cancelled)

63. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 2 and an inert ~~carrier~~carrier.

64. (Cancelled)

65. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 3 and an inert ~~carrier;~~carrier.

66. (Cancelled)

67. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 4 and an inert ~~carrier;~~carrier.

68. (Cancelled)

69. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 10 and an inert ~~carrier;~~carrier.

70. (Cancelled)

71. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 11 and an inert ~~carrier;~~carrier.

72. (Cancelled)

73. (Withdrawn-currently amended) A composition for suppressing transcription of a Type I collagen gene, which comprises a compound according to claim 14 and an inert ~~carrier;~~carrier.